



Field Crops Studies

Volume X

No. 1

2016

***Изследвания
върху полските култури***

***Том X
Книжка 1***

2016

РЕДАКЦИОННА КОЛЕГИЯ:

Гл. РЕДАКТОР: Доц. д-р Юлия Енчева
РЕДАКТОРИ: Проф. д-р Маргарита Нанкова
Проф. д-р Валентина Енчева
Проф. д-р Емил Пенчев
Доц. д-р Татяна Петрова
Доц. д-р Генчо Милев

**ЕЗИКОВИ
РЕДАКТОРИ:** Катя Делчева
Соня Димитрова
гл. ас. д-р Даниела Вълкова

Издател: Добруджански земеделски институт
Редакция: Добруджански земеделски институт
гр. Генерал Тошево, 9520
тел.: +359 58 / 603 125; факс: +359 58 / 603 183
e-mail: fcs@dai-gt.org; <http://fcs.dai-gt.org/>
Корица: Катя Делчева, Стефан Димитров
Дизайн и предпечат: Катя Делчева, Стефан Димитров
Печат: "Нилекта Принт" ООД - гр. Добрич (+359 58 600 299)
ISSN 1312-3882

EDITORIAL BOARD:

EDITOR IN CHIEF: *Assoc. Prof. Julia Encheva*
EDITORS: *Prof. Margarita Nankova*
Prof. Valentina Encheva
Prof. Emli Penchev
Assoc. Prof. Tatyana Petrova
Assoc. Prof. Gencho Milev

**LANGUAGE
EDITORS:** Katia Delcheva
Sonia Dimitrova
Daniela Valkova

Publisher: Dobrudzha Agricultural Institute
Address: Dobrudzha Agricultural Institute
General Toshevo 9520
phone: +359 58 / 603 125; fax: +359 58 / 603 183
e-mail: fcs@dai-gt.org; <http://fcs.dai-gt.org/>
Cover design by Katia Delcheva & Stefan Dimitrov
Text design and typeset by Katia Delcheva & Stefan Dimitrov
Printed by Nilekta Print Ltd. - Dobrich (+359 58 600 299)
ISSN 1312-3882

EFFECTS OF A BIOSTIMULANT AND A MINERAL FERTILIZER ON THE ANTIOXIDATIVE DEFENCE SYSTEM OF CHILLING-EXPOSED MAIZE PLANTS

Rositsa Cholakova-Bimbalova, Lubka Koleva, Andon Vassilev
Agricultural University – Plovdiv

Abstract

Cholakova-Bimbalova, R., L.Koleva, A., Vassilev. Effects of a biostimulant and a mineral fertilizer on the antioxidative defence system of chilling-exposed maize plants. FCS 10(1):207-214

The aim of the study was to investigate the effects of the biostimulant Terra-Sorb Foliar and the mineral fertilizer Poly-Plant on the antioxidative defence system of chilling exposed young maize plants (*Z. mays* L., hybrid Kneza 307). The plants were grown as a substrate-hydroponic culture in a controlled environment at 25±1°C/20±1°C (day/night) temperature. At the appearance of the third leaf, the maize plants were exposed to chilling stress (constant 10°C) for 14 days. Control plants were grown at 25/20°C. Seven days after the beginning of the chilling treatment, a part of the chilled plants were sprayed with 1 % water solution of the biostimulant Terra-Sorb Foliar and with 1 % water solution of the mineral fertilizer Poly-Plant. The plants were left to grow for another 7 days at the same conditions. At the end of the experimental period the antioxidant activity, polyphenols and anthocyanins content as well as guaiacol peroxidase (GPOD) activity were determined. The applied biostimulant and the mineral fertilizer ameliorated to some extent the negative effect of chilling treatment of the young maize plants.

Key words: *Zea mays* L. – chilling – antioxidant activity – anthocyanin – polyphenols – biostimulant – leaf fertilizer

Резюме

Чолакова-Бимбалова, Р., Л. Колева, А., Василев. Изпитване влиянието на биостимулант и минерален тор върху антиоксидантната защитна система на царевични растения, подложени на нискотемпературен стрес. FCS 10(1):207-214

Целта на изследването беше да се проучат ефектите на биостимуланта Terra-Sorb Foliar и минералния тор Poly-Plant върху антиоксидантната защитна система на млади царевични растения (*Z. mays* L., хибрид Кнежа 307), подложени на нискотемпературен стрес. Растенията бяха отглеждани като субстратно-хидропонна култура в контролирана среда при температура 25±1°C/20±1°C (ден/нощ). При появата на трети лист царевичните растения бяха изложени на нискотемпературен стрес (постоянна температура 10°C) за 14 дни. Контролните растения бяха отглеждани при температурен режим 25/20°C. Седем дни след началото на нискотемпературното въздействие, част от охладените растения бяха изпръскани с 1% воден разтвор на биостимуланта Terra-Sorb Foliar и с 1% воден разтвор на минералния тор Poly-Plant. Растенията бяха оставени да се развиват за още 7 дни при същите условия. В края на експерименталния период бяха определени общата антиоксидантна активност, съдържанието на полифеноли и антоциани, както и активността на гваякол

пероксидаза (GPOD). Приложението на биостимуланта и минералния тор намалиха до известна степен отрицателното влияние на ниските положителни температури върху младите царевични растения.

Ключови думи: *Zea mays* L. – ниски положителни температури - антиоксидантна активност - антоциани - полифеноли - биостимулант - листен тор

INTRODUCTION

Chilling is one of the major factors limiting growth, distribution, and productivity of tropical plants, such as maize (*Zea mays* L.). Chilling adversely affects the physiological processes in maize plants (Hund et al. 2007), especially when they are in early stage, during the transitional period from heterotrophic to autotrophic nutrition (Stamp 1984). Temperatures below 15°C may cause chilling stress and induce different physiological disorders in young maize plants (Hola et al. 2007; Leipner 2009).

During chilling stress, the production of reactive oxygen species (ROS) is accelerated, which may damage proteins, nucleic acids, lipids and other molecules (Aroca et al. 2003). Plants have several mechanisms to prevent or alleviate the damage from ROS. The mechanisms include non-enzymatic antioxidants, metabolites such as ascorbate, alfa-tocopherol, etc. and enzymatic antioxidants, such as SOD, CAT, POX, APX, etc. (Farooq et al. 2008; Foyer et al. 1994; Pinhero et al. 1997). Other secondary products of the metabolite processes, which are synthesized very intensively under chilling stress, such as anthocyanins, phenols, etc. possess a radical scavenging activity and act as antioxidants (Christie et al. 1994; Rivero et al. 2001).

Chilling temperatures may provoke decrease of water and mineral uptake (Aroca et al. 2003), stomatal conductance (Melkonian et al. 2004) and photosynthetic activity (Foyer et al. 2002). The negative effects have further consequences (Aroca et al. 2003; Foyer et al. 2002; Melkonian et al. 2004) leading to plant growth retardation (Leipner 2009; Zaidi et al. 2010).

The application of different biostimulants and leaf fertilizers has been used to improve plant tolerance to environmental stresses, such as drought (Petrozza et al. 2014), low temperature (Botta 2013), salinity (Ertani et al. 2013) and others. Surprisingly, there is a very little information about the effects of biostimulants and leaf fertilizers on the physiological status of maize plants exposed to chilling temperatures. Therefore, the aim of our study was to evaluate the effects of a biostimulant and a mineral fertilizer on an antioxidative defence system of young maize plants under chilling stress.

MATERIALS AND METHODS

Growth Conditions and Experimental Design

The experiments were carried out in a climatic room of the Department of Plant Physiology and Biochemistry at the Agricultural University of Plovdiv, Bulgaria. Maize plants, from hybrid "Kneza 307", were grown as a substrate-hydroponic culture, using ½ strength modified Hoagland nutrient solution, at controlled environment: photoperiod – 12 hours, PPFD (Photosynthetic photon flux density) 200 $\mu\text{mol m}^{-2} \text{s}^{-1}$, temperature – 25±1°C/20±1°C (day/night) and relative air humidity – 60±5%. At the appearance of the third leaf, the maize plants were exposed to chilling stress (constant 10°C) for 14 days. Control plants were grown at 25/20°C. Seven days after the beginning of the chilling treatment, a part of the chilled maize plants were sprayed with 1 % water solution of the biostimulant Terra-Sorb Foliar and with 1 % water solution of the mineral fertilizer Poly-Plant. The volume of leaf spraying was 1 ml per plant. The plants were left to grow for another 7 days at the same conditions. The product Terra-Sorb Foliar contains free amino acids and small peptides, while the product Poly-Plant contains macro- and microelements. The experimental design included 4 treatments, namely: control 25°C, chilling 10°C, Terra-Sorb

Foliar 10°C and Poly-Plant 10°C. Each treatment had 3 replications (pots) with 4 plants per pot. Analyses were conducted at the end of the experimental period. The experiment was performed twice.

Antioxidant activity

This parameter was measured with the preformed radical monocation of 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS^{•+}) (Shalaby et al. 2013) in extracts obtained in the way described for total phenolics. The incubation mixture contained 300 µl extract and 3.0 ml ABTS (with 0.7 abs at 734 nm wavelength). After 6 min at room temperature, the absorbance was measured at 734 nm. For reference (blank sample) ABTS + 300 µl 80% MeOH was used. The antiradical activity is expressed as % total antioxidant activity and is equal to:

$$\text{TAA\%}_{(\text{Total antioxidant activity})} = (\text{Abs}_{(\text{plant sample})} - \text{Abs}_{(\text{blank})} / \text{Abs}_{(\text{plant sample})}) * 100$$

Polyphenols content

The total amount of phenolic compounds in the plant extracts was determined with the reagent of Folin-Ciocalteu (Waterman et. al. 1994) following the methodology of Singleton and Rossi (1965), with slight modifications. The samples (1 g of fresh leaf material) were ground with quartz sand and 10 ml 60% acidic methanol, and submerged in an ultrasound bath for 15 min. The homogenized material was then transferred to suitable tubes, which were carefully sealed and left for 15 hours in the dark at room temperature for extraction. During the incubation period, the tubes were periodically stirred. Afterward, the tubes were centrifuged and the supernatant, which was used for the measurement of total phenolics, anthocyanins, and antioxidant activity, was carefully collected in new clean tubes. For the determination of total phenolics were mixed 40 µl of extract, 3160 µl distilled water, 200 µl Folin-Ciocalteu reagent and after a minute was added 600 µl 20% NaCO₃. The test tubes were left for 2 hours at room temperature for the reaction to occur. After that, the extinction at 765 nm wavelength was measured. Total phenolics were calculated as gallic acid equivalents (GAE) using a standard curve and are presented as mg/g fresh weight. The standard curve was prepared with gallic acid (Sigma-Aldrich, St. Louis, MO) in the range 0-500 mg/l.

Anthocyanins content

The measurement of the quantity of monomeric anthocyanins was performed with the pH - differential method (Guisti & Wrolstad 2001). The plant extracts with acidic methanol (as described in the procedure for phenolics measurement) were diluted with a buffer (0.025 mol/l potassium chloride), adjusted to pH=1 with HCl, and another buffer (0.4 mol/l sodium acetate) with pH = 4.5. Each sample was diluted to a certain extent with the first buffer with pH=1 (which gives the value DF) and the absorption was measured at 520 nm and 700 nm. A second aliquot of each sample was diluted to the same extent with the second buffer with pH = 4.5 and again the extinction at 520 nm and 700 nm was measured. To calculate total anthocyanins content, the absorption values were used in the following formula:

$$A = (A_{\lambda_{520}} - A_{\lambda_{700}})_{\text{pH } 1.0} - (A_{\lambda_{520}} - A_{\lambda_{700}})_{\text{pH } 4.5}$$

$$\text{TA} = (A * \text{MW} * \text{DF} * 1000) / \epsilon * 1$$

where: the molar extinction coefficient (ϵ) for cyanidin-3-glucoside = 26900 (M⁻¹ cm⁻¹), the molecular weight (MW) of cyanidin-3-glucoside = 449.2 g/mol and the dilution factor (DF) demonstrates the times dilution of the sample. The results (TA – total anthocyanins) are expressed as mg cyanidin-3-glucoside chloride/100 g fresh plant material used.

GPOD enzyme activity

To obtain the enzyme extract 1 gram of fresh plant material was homogenized in 5 ml ice-cold extraction 0,1 M Tris-HCl (pH 7,8) buffer containing 1 mM DTT and 1mM EDTA. After that, the samples were centrifuged at 13 500 g (4 °C for 10 min). The supernatant was used to determine the enzymatic activity spectrophotometrically on a UV/VIS spectrophotometer Pharo 300, according to the methodology of Mocquot et al. (1996). The activity of guaiacol peroxidase (GPOD) (EC 1.11.1.7) was measured at 436 nm according to Bergmeyer (1974). The reaction mixture in the cuvette contained: 2,3 ml 0,1 M KH_2PO_4 (pH 7,0), 300 μl 5 mM H_2O_2 , 300 μl guaiacol and 100 μl enzyme extract.

Statistical Analysis

Statistical analysis was performed using one-way ANOVA (for $P < 0.05$). Based on ANOVA results, a Duncan test for mean comparison was performed, for a 95% confidence level, to test for significant differences among treatments. In the figures and the tables, different letters (a, b, c and d) express significant differences at the $P < 0.05$.

RESULTS AND DISCUSSION

The young maize plants subjected to 14 days chilling had higher total antioxidant activity (TAA) in both leaves and roots compared to the control plants. The TAA was increased by 134% in the leaves and by 17% in the roots of the chilled maize plants (Table 1). The absence of stress conditions in the control plants kept the value of the total antioxidant activity low.

The application of the biostimulant Terra-Sorb Foliar and the mineral fertilizer Poly-Plant changed the TAA of the chilled maize plants. The TAA values of the plants from these treatments were diminished by 26% - leaves and 25% - roots for Terra-Sorb Foliar-treated chilled plants and 7% - leaves and 27% - roots for Poly-Plant –treated chilled plants, respectively, as compared to untreated chilled plants.

Table 1. Effects of a biostimulant and a mineral fertilizer on total antioxidant activity (%) in leaves and roots of chilling-exposed young maize plants

Treatment	TAA (%)	
	Leaves	Roots
Control 25°C	27.1±0.6 ^c	11.7±0.1 ^b
Chilling 10°C	63.5±0.3 ^a	13.7±1.1 ^a
Terra-Sorb Foliar 10°C	47.3±0.2 ^b	10.3±0.5 ^c
Poly-Plant 10°C	59.2±1.2 ^a	10.0±0.1 ^c

The data presented is an average \pm SD. Different letters (a, b, c and d) following the mean values indicate significant differences at $P < 0.05$.

The chilling induced changes in anthocyanins and polyphenols content (Table 2 and Table 3) of maize plants, which is in accordance with the higher value of total antioxidant activity. The value of anthocyanins content in leaves of chilled plants raised more than twice compared to that of the control plants and the polyphenols content was enhanced by 75% in leaves and by 42% in roots compared to that of the control plants, respectively.

The application of Terra-Sorb Foliar and Poly-Plant on chilled maize plants reduced the appearance of significant changes in the anthocyanins content. The anthocyanins level in the leaves of Terra-Sorb Foliar-treated chilled plants was by 54% lower compared to the untreated chilled plants and by 30% in the leaves of Poly-Plant-treated chilled plants.

The same tendency is observed in polyphenols content. The polyphenols content in the leaves of Terra-Sorb Foliar-treated chilled plants was diminished by 9% and 30% of the leaves and roots, respectively, as compared to the untreated chilled plants. In Poly-Plant-treated chilled plants the respective values were lowered by 11 and 2%.

Table 2. Effects of a biostimulant and a mineral fertilizer on leaf anthocyanins content (mg cyanidin-3-glucoside/100g FW) of chilling-exposed young maize plants

Anthocyanins content	
Treatment	Leaves
Control 25°C	88,1±3.6 ^d
Chilling 10°C	241,5±12.1 ^a
Terra-Sorb Foliar 10°C	111,3±5.8 ^c
Poly-Plant 10°C	168,1±10.2 ^b

The data presented is an average ± SD. Different letters (a, b, c and d) following the mean values indicate significant differences at P<0.05.

Table 3. Effects of a biostimulant and a mineral fertilizer on polyphenols content (mg GAE/g FW) in leaves and roots of chilling-exposed young maize plants

Polyphenols		
Treatment	Leaves	Roots
Control 25°C	6.1±0.8 ^b	4.3±0.5 ^b
Chilling 10°C	10.7±0.2 ^a	6.1±1.5 ^a
Terra-Sorb Foliar 10°C	9.7±1.1 ^a	4.3±0.3 ^b
Poly-Plant 10°C	9.5±1.0 ^a	6.0±0.5 ^a

The data presented is an average ± SD. Different letters (a, b, c and d) following the mean values indicate significant differences at P<0.05.

The applied 14 days chilling treatment (10°C) increased the activity of guaiacol peroxidase (GPOD) in the leaves and roots of young maize plants from the hybrid Kneza 307. The chilled maize plants enhanced their GPOD activity by 26% in leaves and 76% in the roots as compared to the control plants (Table 4).

The leaf application of both Terra-Sorb Foliar and Poly-Plant lowered the GPOD activity in the chilled plants. The GPOD activity of Terra-Sorb-treated chilled plants was lowered by 29% in the leaves and by 34% in the roots. The respective values in Poly-Plant-treated chilled plants were diminished by 21% in both roots and leaves.

Table 4. Effects of a biostimulant and a mineral fertilizer on guaiacol peroxidase activity (U/g FW) in leaves and roots of chilling exposed young maize plants

Guaiacol peroxidase		
Treatment	Leaves	Roots
Control 25°C	1.9±0.3 ^b	10.9 ± 0.6 ^c
Chilling 10°C	2.4±0.4 ^a	19.2±1.6 ^a
Terra-Sorb Foliar 10°C	1.7±0.1 ^b	12.6±0.8 ^c
Poly-Plant 10°C	1.9±0.2 ^b	15.2±1.3 ^b

The data presented is an average ± SD. Different letters (a, b, c and d) following the mean values indicate significant differences at P<0.05.

The detected responses of both enzymatic and non-enzymatic components of the plant antioxidant system are a result of the negative effects of chilling temperature. Considering the effects of Terra-Sorb Foliar and Poly-Plant on the different components of the antioxidative defence system of the chilled maize plants, we may conclude that they had ameliorating influence.

The observed effects of chilling on young maize plants correspond to the results of other authors' researches work, concerning exposure of warm climate plants to chilling temperatures. Sarkar et al. (2009) reported that the total antioxidant activity (ABTS assay) increased in turfgrass species with gradual reduction of temperature. Singh et al. (2014).

also observed that the decrease in temperature enhanced the total phenolic compounds in maize plants. The increased levels of the total phenolic compounds have been observed in chill-stressed tomato and melon (Rivero et al. 2001), pepino plants (Sivaci et al. 2014), in lettuce leaf basil (Kalisz et al. 2016). Enhanced levels of anthocyanins and peroxidases in warm climate plants subjected to chilling stress have been reported by a number of authors. Pietrini et. al. (2002) observed enhanced anthocyanins content in maize plants exposed to low temperatures. Leipner (2009) also found a higher anthocyanins level in chilled-stressed maize plants. Khorshidi & Moafi (2014) and Kang & Saltveit (2002) reported for enhanced activity of guaiacol peroxidase in maize plants exposed to chilling temperatures. Increased activity of peroxidases, including guaiacol peroxidase, has been reported by Lee & Lee (2000) in the response of cucumber to chilling stress.

The leaf application of the biostimulant Terra-Sorb Foliar and the mineral fertilizer Poly-Plant ameliorated the negative effects of chilling temperature. The components they contain may be easily taken from the leaves and therefore may be involved in different metabolic pathways. The biostimulant Terra-Sorb Foliar contain free amino acids and small peptides. According to observations of Matsumiya and Kubo (2011) amino acids and small peptides can be easily absorbed by the leaves. The leaf uptake of mineral nutrients is a common practice in agriculture, especially when the root mineral nutrition is disturbed. In our previous study (Cholakova & Vassilev 2015) we found that chilling diminished the uptake of macroelements in maize plants, consequently, the leaf application of the fertilizer Poly-Plant cause positive effect of these plants.

The observed positive effects of both biostimulant Terra-Sorb Foliar and foliar fertilizer Poly-Plant on chilled maize plants are in a good correspondence with the results of other authors' research work connected with the influence of similar products on stress-exposed plants. For example, Ertani et al. (2013) used biostimulants to ameliorate the negative effect of salt stress on the physiological status of maize plants. Petrozza et al. (2014) also used biostimulants to improved drought resistance of tomato plants and Botta (2013) - cold tolerance of lettuce plants. It was also shown that the application of protein hydrolysates increased nitrogen assimilation and shoot biomass in hydroponically-grown maize (Schiavon et al. 2008) and secondary plant metabolism (Ertani et. al. 2011; Schiavon et al. 2010). Hu et al. (2008) also conducted some analyses to evaluate the effect of foliar fertilization on the growth and mineral nutrient content of maize seedlings under drought and salinity. These reports gave some evidence that leaf-applied amino acids, small peptides, and mineral elements may get into metabolic pathways and support the recovering process in the stressed plants.

In conclusion, the chilling temperatures caused chilling stress and induced changes in the antioxidant defense system of young maize plants. The negative effects have been slightly ameliorated when the plants were sprayed by the biostimulant Terra-Sorb Foliar as well as by the mineral fertilizer Poly-Plant. Further studies are being in progress to clarify the nature of the detected positive effects of the used biostimulant and mineral leaf fertilizer on the physiological status of chilling-exposed maize plants.

REFERENCES

- Aroca, R., Irigoyen, J.J., Sánchez-Díaz, M., 2003.** Drought enhances maize chilling tolerance. II. Photosynthetic traits and protective mechanisms against oxidative stress. *Physiologia Plantarum* 117: 540–549.
- Bergmeyer, HU., 1974.** Methods of enzymatic analysis. Verlag Chemie, Weinheim, Bergstrasse.
- Botta, A., 2013.** Enhancing plant tolerance to temperature stress with amino acids: an approach to their mode of action. *Acta Hort.* 1009, 29–35.
- Cholakova-Bimbalova, R. and Vasilev, A., 2015.** Influence of low temperatures on the growth and macronutrient content in young maize plants. Agricultural University – Plovdiv, Scientific Works, vol. LIX, book 2, Jubilee Scientific Conference with

- International Participation Traditions and Challenges of Agricultural Education, Science and Business
- Christie, P.J., Alfenito, M.R., Walbot, V., 1994.** Impact of low-temperature stress on general phenylpropanoid and anthocyanin pathways: Enhancement of transcript abundance and anthocyanin pigmentation in maize seedlings. *Planta* 194:541–549.
- Ertani, A., Francioso, O., Tugnoli, V., Righi, V., Nardi, S., 2011.** Effect of commercial Lignosulfonate-Humate on *Zea mays* L. metabolism. *J. Agric. Food Chem.* 59, 11940–11948.
- Ertani, A., Schiavon, M., Muscolo, A., and Nardi, S., 2013.** Alfalfa plant-derived biostimulant stimulate short-term growth of salt stressed *Zea mays* L. plants. *Plant Soil* 364: 145–158. doi: 10.1007/s11104-012-1335-z
- Farooq, M., Aziz, T., Basra, S.M.A., Cheema, M.A., Rehamn, H., 2008.** Chilling tolerance in hybrid maize induced by seed priming with salicylic acid. *Journal of Agronomy and Crop Science* 194: 161-. 168.
- Foyer C. H., Decouvrieres P., Kunert K .J. 1994.** Protection against oxygen radicals: an important defence mechanism studied in transgenic plants. *Plant Cell Environ.* 17, 507-523.
- Foyer, C.H., Vanacker, H., Gomez, L.D., Harbinson, J., 2002.** Regulation of photosynthesis and antioxidant metabolism in maize leaves at optimal and chilling temperatures: review. *Plant Physiology and Biochemistry* 40: 659–668
- Guisti, M. M. and Wroldstad, R. E., 2001.** Characterization and measurement of anthocyanins by UV-Visible spectroscopy. *Current Protocols in Food Analytical Chemistry*, F:F1:F1.2.
- Hola D, Kocova M, Rothova O, Wilhelmova N, Benesova M., 2007.** Recovery of maize (*Zea mays* L.) inbreds and hybrids from chilling stress of various duration: photosynthesis and antioxidant enzymes. *Journal of Plant Physiology* 164; 868–877.
- Hu, Y., Burucs, Z., Schmidhalter, U., 2008.** Effect of foliar fertilization application on the growth and mineral nutrient content of maize seedlings under drought and salinity. *Soil Science and Plant Nutrition* 54, 133–141 doi: 10.1111/j.1747-0765.2007.00224.x
- Hund, A., Richner, W., Soldati, A., Fracheboud, Y., Stamp, P., 2007.** Early vigour of maize inbred lines at low temperature as characterised by root and shoot traits. *Eur. J. Agron.* 27: 52–61.
- Kalisz, A., Pokluda, R., Neugebauerová, J., 2016.** Chilling-induced changes in the antioxidant status of basil plants *Acta Physiologiae Plantarum*, DOI: 10.1007/s11738-016-2214-7
- Kang, H.M. and Saltveit, M. E., 2002.** Chilling tolerance of maize, cucumber and rice seedling leaves and roots are differentially affected by salicylic acid. *Physiologia plantarum* 115: 571–576. 2002
- Khorshidi, M., and Moafi, S.J., 2014.** Antioxidative properties in two maize cultivar under low temperature, *International Journal of Farming and Allied Sciences* Available online at www.ijfas.com ©2014 IJFAS Journal-2014-3-6/616-622/ 30 June, 2014 ISSN 2322-4134 ©2014 IJFAS
- Lee, D.H. and Lee, C.B., 2000.** Chilling stress-induced changes of antioxidant enzymes in the leaves of cucumber: in gel enzyme activity assays. *Plant Sci.*, 159 (1), pp. 75-85
- Leipner, J. and Stamp, P., 2009.** Chilling stress in maize seedlings. In: Bennetzen, J.L., Hake, S.C. (eds) *Handbook of Maize: Its Biology*, pp. 291-310. Springer, Heidelberg.
- Matsumiya, Y., Kubo, M., 2011.** Soybean peptide: novel plant growth promoting peptide from soybean. In: El-Shemy, H. (Ed.), *Soybean and Nutrition*. In Tech Europe Publisher, Rijeka, pp. 215–230
- Mocquot, B., Vangronsveld, J., Clijsters, H., Mench. M., 1996.** Copper toxicity in young maize (*Zea mays* L.) plants: Effects on growth, mineral and chlorophyll contents, and enzyme activities. *Plant and Soil* 182: 287–300
- Melkonian J., Yu L. X., Setter T. L., 2004.** Chilling responses of maize (*Zea mays* L.)

- seedlings: root hydraulic conductance, abscisic acid, and stomatal conductance. *J. Exp. Bot.* 55 1751–1760 10.1093/jxb/erh215
- Petrozza, A., Santaniello, A., Summerer, S., Di Tommaso, G., Di Tommaso, D., Paparelli, E., 2014.** Physiological responses to Megafol® treatments in tomato plants under drought stress: a phenomic and molecular approach. *Sci. Hort.* 174: 185–192.
- Pietrini, F., Iannelli, M. A., Massacci, A., 2002.** Anthocyanin accumulation in the illuminated surface of maize leaves enhances protection from photo-inhibitory risks at low temperature, without further limitation to photosynthesis. *Plant, Cell and Environment* 25: 1251–1259
- Pinhero, R.G., Rao, M.V., Paliyath, G., Murr, D.P., Fletcher, R.A., 1997.** Changes in activities of antioxidant enzymes and their relationship to genetic and paclobutrazol-induced chilling tolerance of maize seedlings. *Plant Physiol.* 114:695–704
- Rivero, M., Ruiz, J. M., Garcí'a, P. C., Lo'pez-Lefebvre, L. R., Sa' nchez, E., Romero, L., 2001.** Resistance to cold and heat stress: accumulation of phenolic compounds in tomato and watermelon plants, *Rosa Plant Science* 160: 315–321
- Sarkar, D., Bhowmik, P.C., Young-In-Kwon, Shetty, K., 2009.** Cold Acclimation Responses of Three Cool-season Turfgrasses and the Role of Proline-associated Pentose Phosphate Pathway Dipayan J. AMER. SOC. HORT. SCI. 134(2):210–220.
- Schiavon, M., Ertani, A., Nardi, S., 2008.** Effects of an alfaalfa protein hydrolysate on the gene expression and activity of enzymes of TCA cycle and N metabolism in *Zea mays* L. *J. Agric. Food Chem.* 56, 11800–11808.
- Schiavon, M., Pizzeghello, D., Muscolo, A., Vaccaro, S., Francioso, O., Nardi, S., 2010.** High molecular size humic substances enhance phenylpropanoid metabolism in maize (*Zea mays* L.) *J. Chem. Ecol.*, 36 (2010), pp. 662–669
- Shalaby, E. and Shanab, S.M., 2013.** Comparison of DPPH and ABTS assays for determining antioxidant potential of water and methanol extracts of *Spirulina platensis*. *Indian J. Geo-Marine Sci.* 42 (5): 556-564
- Singh, S.K., Reddy, K.R., Reddy, V.R., Gao, W., 2014.** Maize growth and developmental responses to temperature and ultraviolet-B radiation interaction. *Photosynthetica* 52: 262. doi:10.1007/s11099-014-0029-6
- Singleton, V.L. and Rossi, J.A., 1965.** Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *American Journal of Enology and Viticulture*, 16:144-153.
- Sivaci, A., Kaya, A., Duman, S., 2014.** Effects of ascorbic acid on some physiological changes of pepino (*Solanum muricatum* Ait.) under chilling stress. *Acta Biol Hung.* 65(3): 305-18. doi: 10.1556/ABiol.65.2014.3.7.
- Stamp, P. 1984.** Chilling tolerance of young plants demonstrated on the example of maize (*Zea mays* L.). *J. Agron. Crop Sci.* 7 1–83.
- Waterman P.G. and Mole, S., 1994.** Analysis of phenolic plant metabolites. Oxford, Blackwell Scientific Publications, 238 p
- Zaidi, P.H., Yadav, M., Maniselvan, P., Khan, R., Shadakshari, T.V., Singh, R.P., Pal, D., 2010.** Morpho-physiological traits associated with cold stress tolerance in tropical maize (*Zea mays* L.). *Maydica.* 55: 201-208.