

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

7,200

Open access books available

192,000

International authors and editors

210M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

14%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# Sustainable Pasture Management

---

Atanas Sevov, Chtistina Yancheva and  
Yanka Kazakova

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.72310>

---

## Abstract

Grasslands which are a major part of the global ecosystem, covering 37% of the earth's terrestrial area, have a significant contribution to food security through providing most of the energy and proteins required by the ruminants used for meat and dairy production. Grasslands are considered to have the potential to play a fundamental role in climate change mitigation, particularly regarding carbon storage and sequestration and for biodiversity preservation. This chapter provides an overview of the causes of the pasture degradation and some essential elements for sustainable management, which aims to improve the quantity and quality of pasture, mitigation of climate change and biodiversity preservation. Another point of this chapter is the grasslands with high nature value that nowadays is a top priority in the European legislation as the European Commission has confirmed that HNV farming will remain a key priority in 2014–2020. We present the situation in Bulgaria because it is one of the first member state countries that have assessed HNV regions and put funding in place to support them.

**Keywords:** grasslands, grass composition, perennial grasses, uncontrolled grazing, sustainable management, rotational grazing, high nature value (HNV), Bulgarian grasslands

---

## 1. Introduction

Future challenges related to the sustainable management of natural resources and investments in food production, agriculture and biotechnology research can be summarized as follows: global population growth (the population of the earth will be about 9.2 billion in 2050), global climate change and its adverse impact on agriculture [1], depletion of natural resources with significant importance for the development of world agriculture (e.g. global phosphorus deposits), food safety and security and new ethical requirements for producers.

---

Grasslands which are a major part of the global ecosystem, covering 37% of the earth's terrestrial area, have a significant contribution to food security through providing most of the energy and proteins required by the ruminants used for meat and dairy production. Grasslands are considered to have the potential to play a fundamental role in greenhouse gas (GHG) mitigation, particularly regarding carbon storage and sequestration. Conant et al. [2] conclude that grasslands can act as a significant carbon sink with the implementation of improved management. According to the estimation of FAO [3], global carbon stocks in grasslands is about 343 Gt C, which is about 50% more than the amount stored in world forests.

O'Mara [4] indicates that grazing management and pasture improvement have a global technical potential for mitigation of almost 1.5 Gt carbon dioxide equivalents in 2030, with additional reduction possible from the restoration of degraded lands. According to Nordborg and Rööös [5], the total carbon storage potential in pastures does not exceed 0.8 tons of C per ha and year or 27 billion tons of C globally. During the last years, many researchers studied the function of grasslands as a carbon sink and the main factors affecting the storage process [2, 5–11, 46]. According to some authors [6, 9, 10, 12, 18, 22, 27], soil's grazing intensity (under- and overgrazing) can lower carbon sequestration or lead to carbon losses. These authors observed effects of grazing mediated by changes in the removal, growth, carbon allocation and flora in pastures and carbon input from ruminant excreta, which affect the amount of carbon in soils [27, 36, 40].

The results of the studies conducted by Alemu et al. [14] indicated that grazing management practices impacted greenhouse gas (GHG) intensity of beef production by affecting diet quality, animal performance and soil C change. It also emphasizes the importance of accounting for all emission sources and sinks within a beef production system when estimating its environmental impacts.

## **2. Effect of continuous (uncontrolled) grazing on grasslands**

Formation and development of grass compositions in meadows and pastures were conditioned with the influence of soil and weather, relief, altitude, plant species interactions, micro-organisms, animals and humans. All these factors are interrelated and constantly changing due to variations in the species composition and the quantitative ratio of the different species and groups. Plants in meadows and pastures are changing relatively fast under the influence of different anthropogenic pressures, which can cause both positive and negative changes.

Many high nature value pastures have been abandoned. The meadow mowing has been ceased, which leads to developing of more aggressive grass species, shrubs and trees. Wood and shrub forms begin due to uncontrollable spread or the existence of a forest near the grassland, which gradually spreads from the end to the middle of the area.

Due to the weak animal's grazing efficiency in seminatural grasslands, many of them are degraded and turned into arable land, orchards or vineyards.

This leads to the irreversible loss of diversity of plant species as well as of vertebrate and invertebrate species.

The economic status of meadows and pastures is determined by different characteristics of the grasses in them, while the most important indicator is the lawn productivity in normal climatic conditions, which depends on the soil fertility and regimen of use.

Another indicator of the economic status of meadows and pastures is the quality of green mass and hay, which is determined by the degree of acceptance by the animals, nutritional value (protein, vitamins, mineral, salts) and digestibility of the plant species which are part of the grassland.

Many plant species from different botanical families are found in natural meadows and pastures. In comparatively similar areas, soil and climatic conditions, the number of species in grasslands often exceed 50–60. Meadow and pasture grasses are divided into three groups, cereals, legumes and grasses, from other botanical families, which are referred to as “various plants.”

The widest distribution among grasslands has the species from Poaceae family up to 50–90% of the grass [15, 16]. This is due to their highly competitive ability, longer shelf life and durability of unfavorable climatic and soil conditions. Cereal grasses are wanted component in the grasslands because they supply the animals with easily digestible and rich of nutrition biomass. They also protect the soil from water and wind erosion due to dense grass they develop.

Legumes have the highest nutritional value but are less widespread—very often with 6–10%. Their seldom occurrence in meadows and pastures is due to their greater rigor to the environment and the less durability of most species in the family. Only in conditions very favorable to their development, they can reach 50–60% [15, 16]. Compared to cereals and various plants (from other botanical families), legumes are less common in meadows and pastures. They are not a constant element of grasslands, and their participation is strongly influenced by climatic conditions—in wet years, the so-called clover years are more abundant, and in dry years, their involvement is insignificant. Increasing legumes is a way to improve the quality of grassy biomass, as they are rich in proteins, minerals and vitamins. Their higher number in grasslands improves the nitrogen balance of the soil and promotes the more active development of the other species.

The distribution of the various species (from other botanical families), in grasslands, is determined by the peculiarities of the environment—their participation varies from 10 to 60% [15, 16]. This group is distinguished by a great variety of species—there are about 200, with different nutritional values [15]. This group is represented in mountain meadows and high mountain pastures as well as in wet meadows and pastures.

In the grass cover of meadows and pastures, the perennial grasses of these groups prevail. One-year species rarely occur, with greater participation in degraded grasslands as well as in abandoned orchards. These species are important for early grazing in the southern and southeast parts of Bulgaria [16, 28].

Proper and regulated pasture loading is of paramount importance for ensuring quality grazing with valuable botanical composition, conservation of species diversity and longer use. Still, in many countries, free grazing is applied, which damages both the grasses density and species proportion in grasslands.

Also, soil compaction leads to a change in its physical properties, which affects the development of the grass. Species that require better aeration quickly drop out of grasslands and in their place develop more stable grasses that are less productive and inferior in quality. In this way of grazing, only 40% of the grass is used [45].

Grazing has a strong influence on grass composition. Changes in grassland under the grazing influence depend on the kind of animals, the time and the way of grazing, the soil conditions, the grassland peculiarity, etc. [44]. Grazing early in the spring can suppress some valuable species and allow domination of the weeds [16, 18].

The soil compaction increases the number of rhizomatous grasses that are less sensitive to soil aeration and reduces the participation of the demand in this regard of rhizome and high-growth bunch grasses.

In moderately wet pastures, grazing contributes to consolidation and compaction of the sward, and in damp pastures, trampling can lead to swamping and allow invasion of some weeds, causing poaching. In the dry grasslands, the steady treading leads to shattering a grass cover. The unfavorable influence of treading strongly occurs at unsystematic grazing when the animals move freely and stay for a long time in the pasture. It is stronger when the pasture was used by cattle. The grass composition in grasslands significantly changes with the grazing and species selection from the animals during grazing. Under the influence of grazing some plants, fall off the grassland. During the grazing, the animals eat almost entirely the leaves of tall grasses, which make their recovery difficult, and they are relatively quickly dropped out of the grassland. Low-growth and rosette plants are recovering faster as they retain their basal leaves and take the lead in grassland. They are well preserved in pastures and species with creeping, rooting stems and inflorescences near the soil surface (white clover, knotgrass, etc.).

In grazing, animals prefer certain plants, while others avoid. In the case of unsympathetic grazing, the species that animals avoid form seeds, and the grasses they prefer reduce their vitality and gradually fall out of grass. In rotational grazing, the influence of the animal selection during grazing is almost eliminated, while the rest of the plants were harvested, until they are re-grazed and with the practice of cutting the grass, left after the grazing, is complete in each grazing cycle. In case of free grazing, the influence of the different animals on grassland components is also more pronounced. It was known that the sheep graze the grass shallow, making it difficult to restore the common pasture grasses, but cattle partially plucked up some species during grazing.

This effect was observed in many investigations [4, 19, 20], reduced growth, tiller numbers, plant cover and changes in botanical composition.

Early spring and late autumn grazing reduce the participation of valuable pasture grasses, which have not accumulated enough reserves to overcome early grazing and survive during winter. This grazing leads to an increase in the participation of the first developed annual species that grow up by seeds.



In cattle grazing, hard stools have some adverse effects. The plants below them suffocate and die, and some nitrophilic species develop around them. The larvae of some insects and helminths that cause animal diseases develop in the field. The urine, rich in nitrogen, favors the development of valuable species.

Intensive grazing is depleting the soil, despite the fact that part of the grass-fed nutrients was restored to the soil with animal stools. Soil degradation reduces the participation of valuable pasture grasses which are demanding for the presence of nutrients and increases the involvement of the low-productive, medium-quality, densely tufted grasses.

The negative consequences of nonsystemic, uncontrolled grazing can only overcome by introducing an appropriate grazing regime. Systematic and organized grazing (regular or parcel) would help to preserve species diversity and grass density.

### 3. Grazing management

Grazing management—combining animal, plant, soil and other environmental components and the grazing methods by which the system is managed to achieve specific goals—improved pasture condition, higher forage yields and animal production with ecological concern [16].

The sustainable grazing management includes a proper stocking rate, livestock type and recovery time for grass regrowth after grazing. It is important to consider the effect of grazing management on pasture growth, tiller density, pasture quantity and quality and soil properties. Many factors affect quantity and quality of pastures like farm topography, weather variation among the seasons, botanical composition, herbage cover, stocking rate, seasonal grazing, anti-quality compounds in grasses and application of different practices [23, 24].

Rotational grazing was a component of the institutional and scientific response to severe rangeland degradation at the turn of the twentieth century, and it has since become the professional norm for grazing management [25, 31].

What is rotational grazing— all cases in which only one part of pasture is grazed while all other parts rest? That means that the pasture is divided into a certain number of small areas (paddocks), and the livestock can use only one of them. In this case, the grazing animals are moved from one paddock to another and are thus forced to graze much of the grass. In all other (resting) parts, the grass can renew its energy reserves deepen the root system and in the future time to give a maximum production.

Why use rotational grazing? All over the world, people with livestock and grazing land can benefit from rotational grazing. This has some advantages, called benefits such as economic benefits, time savings, environmental benefits, esthetics and human health benefits, better animal health, etc.

A rotational grazing system is preferred in pasture-based animal production because meat from cows and lambs has better quality with less fat, more vitamin E [26, 29] and higher levels of omega-3 and conjugated linoleic fatty acids than grain-finished products [17, 21, 30].

The fundamental advantage for the animals in grazing systems is that the livestock in pastures is healthier than those housed in confinement. The key of sufficient rotational grazing is determination of the number and size of paddocks, water supply for livestock and fence type. Determination of the suitable number of paddocks depends on the time required for grass regrowth (**Table 1**) and grazing period that is varied from 4 to 6 days.

Many authors in their publications present different grazing management models with the consideration of periods of strong growth and animal pressure [7, 43]. Many farmers in countries with hill pastures applied adopted regime with 3 days per paddock and high stocking rate [23].

Jacobo [32] observed that productivity and sustainability might be compatible by replacing continuous with rotational grazing. The reason is that rotational grazing promoted functional groups composed of high forage value species and reduced bare soil through the accumulation of plant residues. These changes indicate an improvement in rangeland condition and in carrying capacity.

These results are relevant to the other authors. As an example, Pavlů et al. [13] studied continuous stocking and rotational grazing. On the base of the databases, authors conclude that vegetation varied as a result of time and differences between treatments. Several prostrate dicotyledonous species (*Trifolium repens* L., *Taraxacum* sp., *Bellis perennis* and *Leontodon autumnalis*) increased under continuous stocking. This treatment also promoted the growth of the perennial grass *Lolium perenne* L., which was able to cope with frequent defoliation. Tall grasses sensitive to frequent defoliation (*Poa trivialis* L., *Holcus mollis* L., *Alopecurus pratensis* L., *Dactylis glomerata* L. and *Elytrigia repens* L.) were more abundant in rotationally grazed paddocks. Species diversity was not significantly influenced by the different grazing systems. The decrease in the potential sward height under continuous stocking revealed the replacing of tall dominants by lower species. Information about pasture management should, therefore, involve not only grazing intensity but also the grazing system used.

The new opportunity to improve the management and welfare of extensively produced beef cattle is to combine technologies for monitoring the spatial behavior of livestock with technologies that monitor pasture availability. According to Manning et al. [33], the Global Navigation Satellite System (GNSS) technology could determine livestock grazing preference and hence improve management and paddock utilization. The cattle behavior changed, highlighting how technologies that monitor these two variables may be used in the future as management tools to assist producers better manage cattle and to manipulate grazing intensity and paddock utilization.

Species	Cool weather	Hot weather
Cool season grasses	14	35–50
Warm season grasses	35–40	21
Legumes	21–28	21–28

Source: Blanchet et al. [31].

**Table 1.** Optimal rest period for forage species in days.

Sustainable pasture management including the application of new technologies has several environmental advantages over tilled land—significantly decrease soil erosion, require minimal pesticides and fertilizer usage and reduce the amount of barnyard runoff. This leads to the conclusion that, taking advantage of wildlife, we can also increase the pasture productivity.

#### 4. High nature value (HNV) grasslands in Bulgaria

By definition high nature value (HNV) farmland represents areas where “agriculture is a major (usually the dominant) land use and where that agriculture supports, or is associated with, either a high species and habitat diversity or the presence of species of European, and/or national, and/or regional conservation concern, or both” [34, 35]. The majority of HNV farmland in Europe comprises seminatural pastures, meadows and orchards as well as various landscape elements [35, 38]. Around one-third of the agricultural area in Bulgaria is potentially of high nature value, and the most significant share of it is seminatural pastures and meadows [41, 45]. The figures below visualize high nature value grasslands in Bulgaria: flower-rich meadows in Elena municipality (**Figure 1**), species-rich pastures in Central Balkans (**Figure 2**) and species-rich pastures in Eastern Stara Planina (**Figure 3**).

HNV grasslands are of particular importance for nature conservation and the European ecological network of protected areas of Natura 2000. There are 18 habitats of natural and seminatural grassland ecosystems in the Bulgarian Natura 2000 sites, which cover between 15 and 20% of their territory [37].

Key features of the HNV farming systems are the low inputs, low outputs and high labor requirements usually resulting in a significant number of species and structural diversity in space and time [38]. The practices most often associated with HNV pastures and meadows are extensive grazing as presented on **Figure 4** and cutting hay (mowing) once or twice per year. **Figure 5** shows traditional hay storage still preserved in Western Stara Planina.



**Figure 1.** Flower-rich meadows in Elena municipality (June 2012, Y. Kazakova).





**Figure 2.** Species-rich pastures in Central Balkans (July 2016, Y. Kazakova).



**Figure 3.** Species-rich pastures in Eastern Stara Planina (June 2012, Y. Kazakova).

However, the modernization of agriculture inevitably leads to the intensification of the traditional practices and decrease in the high nature value. For example, over 90% of the grassland habitats in the European ecological network Natura 2000 are in unfavorable conservation



**Figure 4.** Extensive sheep grazing in Central Stara Planina (July 2016, Y. Kazakova).



**Figure 5.** Traditional hay storage still preserved in Western Stara Planina (November 2014, Y. Kazakova).

status [37]. The two extreme examples are the loss of HNV grasslands due to conversion to intensive meadows or even arable land and the abandonment of farming in areas unsuitable for intensification. **Figure 6** presents scrub overgrowth and closure of landscapes in abandoned grasslands in Western Stara Planina.

The trend is best revealed by the statistical data on grasslands in Bulgaria, presented in **Figure 7** (BANCIK MAF, 2000–2016). The total area of grasslands in Natura 2000 was just over 1.8 million hectares. In 2015, it was down to 1.36 million hectares, a decrease of 24% [41].

Overall, grasslands cover around one-third of the agricultural area in Bulgaria. In the agricultural land use surveys, they are divided into four grassland groups:

1. Permanent productive meadows, which can be natural or planted for longer than 6 years and can be used either for mowing or for grazing. Their area decreased by 38% from 2000 to 2015. Due to their high productivity, they are often converted to arable land.



**Figure 6.** Grasslands abandonment leads to scrub overgrowth and closure of landscapes in Western Stara Planina (April 2015, Y. Kazakova).






Figure 7. Total area of grasslands in Bulgaria (2000–2015).

2. High mountain pastures are located at altitudes between 1000 and 1500 m a.s.l. and are used for summer grazing of livestock. Their area is most stable in comparison to other grassland groups—it decreased by only 6% from 2000 to 2015.
3. Low productivity grasslands—usually used for grazing.
4. Due to their low productivity, they are never mown. They decreased by 19% from 2000 to 2015 mostly due to an abandonment of farming (Figure 7).
5. Orchard meadows, which are permanent productive pastures in orchards with less than 100 trees per hectare. Their area decreased the most, by 46%, from 2000 to 2015.

Another negative tendency for the loss of HNV grasslands is their sale for development. The extensive land use and the species-rich grasslands, as well as the site's characteristics, often create landscapes that are attractive for tourists as shown in Figure 8. This creates development pressure, and the values that attracted visitors ultimately were lost.



Figure 8. Grasslands for sale: the attractive high nature value landscapes stimulate tourism development (June 2012, Y. Kazakova).

When HNV farmlands were first identified in 2007 in Bulgaria, the area of HNV grasslands was estimated at 951,256 ha [39]. Only 5 years later, in 2012, the HNV grassland area decreased to 809,530 ha [37]. Even if there were some methodological differences, the decreasing trend is unquestionable.

To preserve and maintain grassland areas of high nature value and the associated species, measures financed under the Bulgarian Rural Development Programme (2014–2020) [42] are being undertaken to promote or restore traditional management practices for seminatural grassland, as follows:

- Keeping the density of livestock units at 0.3–1 LU/ha according to the natural, climatic and soil conditions to ensure the good ecological status of meadows and pastures and maintenance of a permanent grass cover.
- A ban on the use of mineral fertilizers and pesticides.
- Cleaning of undesirable grass and shrub vegetation.
- Consecutive grazing.

Overall, HNV grasslands in Bulgaria require targeted policy support and improved management both from agricultural and conservation point of view to improve the current situation where the forage resources are decreasing because of the loss of grassland area; the natural quality of the remaining grasslands is also declining due to the intensification or abandonment of extensive, low-input practices.

## Author details

Atanas Sevov<sup>1\*</sup>, Chtistina Yancheva<sup>1</sup> and Yanka Kazakova<sup>2</sup>

\*Address all correspondence to: [asevov@yahoo.com](mailto:asevov@yahoo.com)

1 Agricultural University, Plovdiv, Bulgaria

2 University of National and World Economy, Sofia, Bulgaria

## References

- [1] Scollan ND, Enser M, Richardson RI, Wood JD. Effect of forage legumes on the fatty acid composition of beef. *The Proceedings of the Nutrition Society*. 2002;**61**(3A):99A
- [2] Conant RT, Paustian K, Elliott ET. Grassland management and conversion into grassland: Effects on soil carbon. *Ecological Applications*. 2001;**11**(2):343-355
- [3] FAO. Challenges and opportunities for carbon sequestration in grassland systems—A technical report on grassland management and climate change mitigation. *Integrated Crop Management*. 2010;**9**

- [4] O'Mara FP. The role of grasslands in food security and climate change. *Annals of Botany*. 2012;**110**(6):1263-1270. DOI: 10.1093/aob/mcs209
- [5] Nordborg M, Rööös E. Holistic management—A critical review of Allan Savory's grazing method. SLU, Swedish University of Agricultural Sciences & Chalmers. 2016; ISBN: 978-91-576-9424-9
- [6] Deng L et al. Effects of grazing exclusion on carbon sequestration in China's grassland. *Earth-Science Reviews*. 2017;**173**:84-95. DOI: 10.1016/j.earscirev.2017.08.008
- [7] Sheath GW, Carlson WT. Impact of cattle treading on hill land 1. Soil damage patterns and pasture status. *New Zealand Journal of Agricultural Research*. 1998;**41**:271-278
- [8] Schonbach P, Wan HW, Gierus M, Bai YF, Muller K, Lin LJ, Susenbeth A, Taube F. Grassland responses to grazing: Effects of grazing intensity and management system in an inner Mongolian steppe ecosystem. *Plant and Soil*. 2011;**340**(1-2):103-115
- [9] Wang S, Wilkes A, Zhang Z, Chang X, Lang R, Wang Y, Niu H. Management and land use change effects on soil carbon in northern China's grasslands: A synthesis. *Agriculture, Ecosystems & Environment*. 2011;**142**(3-4):329-340. DOI: 10.1016/j.agee.2011.06.002
- [10] Wang D, Wu GL, Zhu YJ, Shi ZH. Grazing exclusion effects on above- and below-ground C and N pools of typical grassland on the Loess Plateau (China). *Catena*. 2014;**123**:113-120. DOI: 10.1016/j.catena.2014.07.018
- [11] Zavaleta ES, Shaw MR, Chiariello NR, Mooney HA, Field CB. Additive effects of simulated climate changes, elevated CO<sub>2</sub>, and nitrogen deposition on grassland diversity. *Proceedings of the National Academy of Sciences*. 2003;**100**(13):7650-7654
- [12] Liebzig MA, Morgan JA, Reeder JD, Ellert BH, Gollany HT, Schuman GE. Greenhouse gas contributions and mitigation potential of agricultural practices in northwestern USA and western Canada. *Soil & Tillage Research*. 2005;**83**:25-52. DOI: 10.1016/j.still.2005.02.008
- [13] Pavlů V, Hejcman M, Pavlů L, Gaisler J. Effect of rotational and continuous grazing on vegetation of an upland grassland in the Jizerské Hory Mts., Czech Republic. *Folia Geobotanica*. 2003;**38**(1):21-34
- [14] Alemu AW et al. Assessment of grazing management on farm greenhouse gas intensity of beef production systems in the *Canadian prairies* using life cycle assessment. *Agricultural Systems*. 2017;**158**:1-13. DOI: 10.1016/j.agry.2017.08.003
- [15] Ministry of Environment, Ministry of Agriculture. Sustainable land management. Ministry of Environment—Bulgarian Government Publisher; 2007
- [16] Yancheva C, Angelova S, Koeva R. Management of steppe habitats in pastures of the region of Kaliakra reserve in Bulgaria. In: XIX-th General Meeting of EGF in Multifunction Grasslands; La Roshelle, France. 2002. pp. 860-862
- [17] Yang A, Lanari MC, Brewster M, Tume RK. Lipid stability and meat colour of beef from pasture- and grain-fed cattle with or without vitamin E supplement. *Meat Science*. 2002; **60**(1):41-50. DOI: 10.1016/S0309-1740(01)00103-6



- [18] Wang KB, Deng L, Ren ZP, Li JP, Shangguan ZP. Grazing exclusion significantly improves grassland ecosystem C and N pools in a desert steppe of China. *Catena*. 2016;**137**:441-448. DOI: 10.1016/j.catena.2015.10.018
- [19] Garden DL, Lodge GM, Friend DA, Dowling PM, Orchard BA. Effects of grazing management on botanical composition of native grass-based pastures in temperate south-east Australia. *Australian Journal of Experimental Agriculture*. 2000;**40**:225-245. DOI: 10.1071/EA98010
- [20] Garden DL, Dowling PM, Eddy DA, Nicol HI. The influence of climate, soil, and management on the composition of native grass pastures on the central, southern, and Monaro tablelands of New South Wales. *Australian Journal of Agricultural Research*. 2001;**52**(9):925-936. DOI: 10.1071/AR98184
- [21] Ripoll G, Albertí P, Casasús I, Blanco M. Instrumental meat quality of veal calves reared under three management systems and color evolution of meat stored in three packaging systems. *Meat Science*. 2013;**93**:336-343. DOI: 10.1016/j.meatsci.2012.09.012
- [22] Shaw MR, Zavaleta ES, Chiariello NR, Cleland EE, Mooney HA, Field CB. Grassland responses to global environmental changes suppressed by elevated CO<sub>2</sub>. *Science*. 2002;**298**(5600):1987-1990
- [23] King WM, Rennie GM, Devantier B, Hoogendoorn CJ. Impacts of grazing management on hill country pastures: Principles and practices. *Hill Country – Grassland Research and Practice*. 2016;**16**:203-212
- [24] Lambert MG, Clark DA, Litherland AJ. Advances in pasture management for animal productivity and health. *New Zealand Veterinary Journal*. 2004;**52**:311-319. DOI: 10.1080/00480169.2004.36447
- [25] Briske DD, Derner JD, Brown JR, Fuhlendorf SD, Teague WR, Havstad KM, Willms WD. Rotational grazing on rangelands: Reconciliation of perception and experimental evidence. *Rangeland Ecology & Management*. 2008;**61**(1):3-17. DOI: 10.2111/06-159R.1
- [26] Descalzo AM, Insani EM, Biolatto A, Sancho AM, Garcia PT, Pensel NA. Influence of pasture or grain-based diets supplemented with vitamin E on antioxidant/oxidative balance of argentine beef. *Meat Science*. 2005;**70**:35-44. DOI: 10.1016/j.meatsci.2004.11.018
- [27] Rice CW, Owensby CE. The effects of fire and grazing on soil carbon in rangelands. The potential of US grazing lands to sequester carbon and mitigate the greenhouse effect. 2001. DOI: 10.1201/9781420032468.ch13
- [28] Yancheva C, Angelova S, Guteva Ya, Mihovski Tz. Steppe habitats in the pastures of the region of Kaliakra reserve and their preservation. *Journal of Mountain Agriculture on the Balkans*. 2007:104-109
- [29] Fraser TJ, Rowarth JS. Legumes, herbs or grass for lamb performance. *Proceedings—NZ Grassland Association*. 1996;**58**:49-52
- [30] Gutierrez-Pena R et al. Carcass composition and meat quality of pasture-raised Mallorquina sheep in Balearic Islands. Grassland resources for extensive farming systems in marginal lands: Major drivers and future scenarios. In: *Proceedings of the 19th Symposium of the European Grassland Federation*; 7-10 May 2017; Alghero, Italy. CNR-ISPAAAM; 2017

- [31] Blanchet K, Moechnig H, DeJong-Hughes J. Grazing systems planning guide. University of Minnesota Extension Service; 2000
- [32] Jacobo EJ, Rodríguez AM, Bartoloni N, Deregibus VA. Rotational grazing effects on rangeland vegetation at a farm scale. *Rangeland Ecology & Management*. 2006;**59**(3):249-257. DOI: 10.2111/05-129R1.1
- [33] Manning J et al. The behavioural responses of beef cattle (*Bos taurus*) to declining pasture availability and the use of GNSS technology to determine grazing preference. *Agriculture*. 2017;**7**(5):45. DOI: 10.3390/agriculture7050045
- [34] Andersen E, Baldock D, Bennet H, Beaufoy G, Bignal E, Brower F, Elbersen B, Eiden G, Godeschalk F, Jones G, McCracken DI, Nieuwenhuizen W, van Eupen M, Hennekes S, Zervas G. Developing a High Nature Value Farming Area Indicator. Consultancy Report to the EEA. Copenhagen: European Environment Agency; 2003
- [35] Beaufoy G, Jones G, Oppermann R. High Nature Value Farming in Europe. 35 European Countries – Experiences and Perspectives. Ubstadt-Weiher: Verlag Regionalkultur; 2012
- [36] Soussana JF, Tallec T, Blanfort V. Mitigating the greenhouse gas balance of ruminant production systems through carbon sequestration in grasslands. *Animal*. 2010;**4**:334-350. DOI: 10.1017/S1751731109990784
- [37] Ministry of Environment. National Priority Action Framework for Natura 2000 in Bulgaria for the 2014-2020 period. Ministry of Agriculture—Bulgarian Government Publisher; 2014
- [38] Oppermann R, Paracchini ML. HNV Farming—Central to European Cultural Landscapes and Biodiversity. 35 European Countries—Experiences and Perspectives. Ubstadt-Weiher: Verlag Regionalkultur; 2012
- [39] Ministry of Agriculture. Bulgarian Rural Development Programme for the 2007-2013 Programming Period. Ministry of Agriculture—Bulgarian Government Publisher; 2007
- [40] Liu S, Schleuss PM, Kuzyakov Y. Carbon and nitrogen losses from soil depend on degradation of Tibetan Kobresia pastures. *Land Degradation and Development*. 2016;**28**(4):1253-1262. DOI: 10.1002/ldr.2522
- [41] Ministry of Agriculture. 2000-2016 Agristatistics Annual BANCİK Reports on the Land Cover and Land Use in Bulgaria. Ministry of Agriculture—Bulgarian Government Publisher; 2016
- [42] Ministry of Agriculture. Bulgarian Rural Development Programme for the 2014-2020 programming period. Ministry of Agriculture—Bulgarian Government Publisher; 2014
- [43] Pande TN, Valentine I, Betteridge K, MacKay A, Horne D. Pasture damage and regrowth from cattle treading. *Proceedings of the New Zealand Grassland Association*. 2000;**62**: 155-160
- [44] Shakhane LM et al. Changes in botanical composition on three farmlets subjected to different pasture and grazing management strategies. *Animal Production Science*. 2013; **53**(8):670-684

- [45] Stefanova V, Kazakova Y. HNV Farming in Bulgaria. High Nature Value Farming in Europe. 35 European Countries—Experiences and Perspectives. Ministry of Agriculture - Bulgarian government publisher; 2012
- [46] Zhou ZY, Li FR, Chen SK, Zhang HR, Li GD. Dynamics of vegetation and soil carbon and nitrogen accumulation over 26 years under controlled grazing in a desert shrubland. *Plant and Soil*. 2011;**341**(1-2):257-268

IntechOpen

IntechOpen

